

Impact des troubles de la nutrition et de la déglutition dans les infections

Dr F.Behar

UTNc Hôpital DUPUYTREN, Draveil

- Unité Transversale de Nutrition clinique
- Hôpital DUPUYTREN (Draveil), groupe hospitalier Henri MONDOR
- Créée en 2008
- Selon le PNNS 2006, sur appel à projet de la DHOS en 2006
- 7 UTNc au plan national à la création, environ 57 en 2016
- Seule UTNc gériatrique en 2008, ? en 2016
- Activité de consultations, d'HDJ, de formations aux professionnels, de prise en charge nutritionnelle des patients hospitalisés, de vigilance nutritionnelle au sein de l'hôpital...
- 2 diététiciennes, 1 secrétaire, 1 médecin

Fréquente :

4 à 10 % à domicile


15 à 40 % en institution (maisons de retraites...)

30 à 70 % à l'hôpital



Fréquente :

Population (n=)	% dénutris	Référence
Chirurgie (131)	54	Bistrian et al.1974
Médecine (201)	45	Bistrian et al.1976
Chirurgie (105)	12-55	Hill et al.1977
Médecine (134)	46	Weinsier et al.1979
Hôpital général (200)	31.5	Willard et al.1980
Orthopédie (82)	48	Dreblow et al.1981
Hémodialyse (58)	62	Thunberg et al.1981
Chirurgie (90)	30-40	Perrot et al.1982
Gériatrie (59)	61	Bienia et al.1982
Pédiatrie (515)	30	Girardet et al.1989
Gériatrie (324)	41	Constans et al.1992
Gériatrie (40)	46	Alix et al.1992

- La dénutrition provoque une diminution de la masse musculaire (sarcopénie)





■ La dénutrition provoque une diminution de la force musculaire




- Gros...
- Maigre...
- Ou IMC identique ...
... et composition corporelle différente ...

IMC identique, surface corporelle identique,
...mais composition corporelle différente...




	IMC	Surface Corporelle	Surface muscle	Surface tissu adipeux viscéral	Surface tissu adipeux sous-cutané	Densité
Patient A	25,5	1,8	131	208	225	36
Patient B	25,9	1,9	190	76	80	47,1

Augmentation de la mortalité



SL Lim, KCB Ong, YH Chan, WC Loke, M Ferguson, L Daniels


Figure 2. Cumulative survival in well-nourished and malnourished patients (n= 818).
HR = Hazard ratio.




¹Survival and mortality data from Singapore Death Registry.

²Assessment with Subjective Global Assessment within 48 hours of hospital admission.

³Adjusted for ethnicity, age and gender.



Pourcentage de malades avec infection nosocomiale chez des patients hospitalisés



Schneider S.M. et coll., " Malnutrition is an independent factor associated with nosocomial infections ",
British Journal of Nutrition, 92 : 105-111, 2004

The impact of malnutrition on morbidity, mortality, length of hospital stay and costs evaluated through a multivariate model analysis

M. ISABEL T. D. CORREIA,* DAN L. WAITZBERG†




Fig. 1 Nutritional status at hospital admission, 709 patients.

Table 1 Nutritional status vs incidence of complications

Nutritional status	Complications	Without complications
Well nourished	78 (16.8%)	385 (83.2%)
Malnourished (moderately and severely)	64 (27.0%)*	173 (73.0%)
Only the severely malnourished	24 (42.8%)†	32 (57.2%)

* RR = 1.60 (IC = 1.20–2.14; $P < 0.01$).

† RR = 2.54 (IC = 1.77–3.66; $P < 0.01$).

Influence de la dénutrition sur les complications infectieuses

Clinical Nutrition (2003) **22**(3): 235-239
© 2003 Elsevier Science Ltd. All rights reserved.
doi:10.1016/S0261-5614(02)00215-7

The impact of malnutrition on morbidity, mortality, length of hospital stay and costs evaluated through a multivariate model analysis

M. ISABEL T. D. CORREIA,* DAN L. WAITZBERG†

Table 2 Incidence of infectious complications vs the nutritional status

Infectious complications	Well nourished	Malnourished (moderately and severely)
Pulmonary infection	19** (4.1%)	17** (5.9%)
Urinary infection	9** (1.9%)	9** (3.7%)
Wound infection	10** (2.1%)	4** (1.7%)
Sepsis	5* (1.1%)	9* (3.7%)
Intrabdominal abscess	2* (0.4%)	5* (2.1%)
Extraperitoneal abscess	1** (0.2%)	2** (0.8%)
Septic coagulopathy	1** (0.2%)	1** (0.4%)
Total	47* (10.1%)	47* (19.4%)

* $P < 0.01$; ** NS.

Relations between undernutrition and nosocomial infections in elderly patients

ELENA PAILLAUD¹, STEPHANE HERBAUD¹, PHILIPPE CAILLET¹, JEAN-LOUIS LEJONC¹, BERNARD CAMPILLO¹, PHUONG-NHI BORIES²

Table 3. Anthropometric variables and energy intake in the three groups of patients

<i>n</i>	No infection 116	One infection 38	>One infection 31	<i>P</i>
Weight (kg)	61.2 ± 1.5	58.9 ± 3.1	51.3 ± 1.7	0.0079
BMI (kg/m ²)	23.8 ± 0.5	24.0 ± 1.2	21.2 ± 0.7	0.046
MAC (cm)	27.1 ± 0.4	26.7 ± 1.0	24.1 ± 0.7	0.011
TST (mm)	11.6 ± 0.5	12.9 ± 1.3	9.5 ± 0.8	0.064
BST (mm)	4.9 ± 0.3	5.9 ± 0.9	3.1 ± 0.3	0.011
Energy intake (kcal/day)	1717 ± 40	1474 ± 91	1284 ± 74	<0.0001
Energy intake (kcal/kg/day)	29.3 ± 0.8	27.2 ± 2.1	25.1 ± 1.4	0.089


BMI, body mass index; MAC, mid-arm circumference; TST, tricipital skinfold thickness; BST, bicipital skinfold thickness.

The three groups were compared by ANOVA.

Table 4. Biological variables in the three groups of patients

<i>n</i>	No infection 116	One infection 38	>One infection 31	<i>P</i>
Albumin (g/l)	36.3 ± 0.4	34.6 ± 0.6	31.7 ± 0.9	<0.0001
Transferrin (g/l)	0.26 ± 0.01	0.23 ± 0.01	0.19 ± 0.01	<0.0001
Orosomucoid (g/l)	2.1 ± 0.7	1.5 ± 0.1	1.6 ± 0.1	0.861
CRP (mg/l)	16 ± 2	37 ± 10	44 ± 8	0.0003
Lymphocytes (g/l)	1.7 ± 0.1	1.8 ± 0.3	1.5 ± 0.1	0.438
PMN leukocytes (g/l)	4.4 ± 0.2	6.1 ± 1.0	6.7 ± 0.7	0.0027


Le système immunitaire



CPA : cellule présentatrice d'antigène
 LB : lymphocyte B
 LT8 : lymphocyte T CD8
 LT4 : lymphocyte T CD4
 P : plasmocyte


PréTc : lymphocyte T précytotoxique
 LTPC : lymphocyte T cytotoxique
 m : mémoire
 IgC : anticorps
 M : macrophage

La dénutrition perturbe le système immunitaire




Alimentation dépourvue en taurine
chez le chat: aucune phagocytose,
aucune explosion oxydative


La dénutrition perturbe le système immunitaire




La dénutrition perturbe le système immunitaire



La dénutrition perturbe le système immunitaire



La dénutrition perturbe le système immunitaire



D

La vitamine D

C

B

A

0

Pourquoi les sanatoriums avaient-ils une architecture héliotropique ?

30

francetvpluzz

Rations quotidiennes pour couvrir les besoins

1 c-à-café et $\frac{1}{2}$ Huile de foie de Morue

720 g de Girolles

120 g de Harengs au vinaigre


20 Sardines à l'huile

22 Œufs durs


5 kg de Foie de veau

1,250 g de Beurre


La dénutrition perturbe le système immunitaire




La dénutrition perturbe le système immunitaire




La dénutrition perturbe le système immunitaire




Influence de la dénutrition sur le système immunitaire






Influence des infections sur la dénutrition




Influence des infections sur la dénutrition



Moins d'apports

Plus de dépenses

Influence des infections sur la dénutrition



Impact sur la dépense énergétique de repos






Figure 7.3. Based on data of Kinney et al. (46). Reproduced from Elwyn DH: Nutritional requirements of surgical patients. *Crit Care Med* 8:9-20, 1980. With permission of *Critical Care Medicine*. The effects of injury, sepsis, and nutritional depletion on resting energy expenditure.


Le syndrome inflammatoire provoque une dénutrition




Le syndrome inflammatoire provoque une dénutrition




Le syndrome inflammatoire provoque une dénutrition




Le syndrome inflammatoire provoque une dénutrition




Le syndrome inflammatoire provoque une dénutrition





Le syndrome inflammatoire provoque une dénutrition




Le syndrome inflammatoire provoque une dénutrition









Conséquences médico-économiques



Age and Ageing 2005; **34**: 619–625
doi:10.1093/ageing/af1197

Relations between undernutrition and nosocomial infections in elderly patients

ELENA PAILLAUD¹, STEPHANE HERBAUD¹, PHILIPPE CAILLET¹, JEAN-LOUIS LEJONC¹, BERNARD CAMPILLO¹, PHUONG-NHI BORIES²

Table 1. Characteristics of the patients

<i>n</i>	No infection 116	One infection 38	>One infection 31	<i>P</i>
Age (years)	80.0 ± 0.7	84.4 ± 1.2	85.2 ± 1.3	0.0002
Gender (M/F) (<i>n</i>)	37/79	3/35	5/26	0.006
MMS	23.7 ± 0.7	20.1 ± 1.4	18.1 ± 1.5	0.0005
Length of stay (days)	60.1 ± 4.9	78.9 ± 10.7	122.2 ± 19.5	<0.0001
Time of infection after admission (days)		44.7 ± 10.4	22.0 ± 3.8	0.0293

Table 5. Association of clinical characteristics with infection

<i>n</i>	No infection 116	One infection 38	>One infection 31	<i>P</i>
Outcome				<0.0001
Nursing home	18 15.4%	9 22.5%	13 40.6%	
Return home	90 76.9%	22 55.0%	10 31.3%	
Death	5 4.3%	8 20.0%	9 28.1%	
Hospitalisation	4 3.4%	1 2.5%	0 0%	

Conséquences médico-économiques

SL Lim, KCB Ong, YH Chan, WC Loke, M Ferguson, L Daniels

Malnutrition and its impact on cost of hospitalization, length of stay, readmission and 3-year mortality

Comparison of hospitalization outcomes, costs and mortality between well-nourished and malnourished patients (n=818).

Outcome variables	Well-nourished (n=583)	Malnourished ^a (n=235)
Length of hospital stay (days), mean $\pm$ sd	4.6 $\pm$ 5.6	6.9 $\pm$ 7.3
Length of hospital stay (days), median (range)	3 (1-63)	4 (1-59)

Conséquences médico-économiques

SL Lim, KCB Ong, YH Chan, WC Loke, M Ferguson, L Daniels

Malnutrition and its impact on cost of hospitalization, length of stay, readmission and 3-year mortality

Comparison of hospitalization outcomes, costs and mortality between well-nourished and malnourished patients (n=818).

Outcome variables	Well-nourished (n=583)	Malnourished ^a (n=235)
Readmission within 15 days of index admission, n (%)	61 (10.5)	40 (17.0)
Readmission within 90 days of index admission, n (%)	133 (22.8)	87 (37.0)
Readmission within 6 months of index admission, n (%)	187 (32.1)	113 (48.1)

Conséquences médico-économiques

SL Lim, KCB Ong, YH Chan, WC Loke, M Ferguson, L Daniels

Malnutrition and its impact on cost of hospitalization, length of stay, readmission and 3-year mortality

Comparison of hospitalization outcomes, costs and mortality between well-nourished and malnourished patients (n=818).

Outcome variables	Well-nourished (n=583)	Malnourished ^a (n=235)
Inpatient mortality, n (%)	2 (0.3)	10 (4.3)
1-year mortality, cumulative, n (%)	24 (4.1)	80 (34.0)
2-year mortality, cumulative, n (%)	39 (6.7)	100 (42.6)
3-year mortality, cumulative, n (%)	58 (9.9)	114 (48.5)

Conséquences médico-économiques

SL Lim, KCB Ong, YH Chan, WC Loke, M Ferguson, L Daniels

Malnutrition and its impact on cost of hospitalization, length of stay, readmission and 3-year mortality

Comparison of hospitalization outcomes, costs and mortality between well-nourished and malnourished patients (n=818).

Outcome variables	Well-nourished (n=583)	Malnourished ^a (n=235)
Cost of hospitalization for individual patient (S\$), mean $\pm$ sd	3707 $\pm$ 5541	4606 $\pm$ 6665
Cost of hospitalization for individual patient (S\$), median (range)	1897 (179-70472)	2534 (297- 50215)

0148-6071/05/2901-0S57\$03.00/0

JOURNAL OF PARENTERAL AND ENTERAL NUTRITION

Copyright © 2005 by the American Society for Parenteral and Enteral Nutrition

Preoperative Immunonutrition: Cost-Benefit Analysis

Marco Braga, MD*; and Luca Gianotti, MD, ScD†

Evaluation économique d'un traitement par Impact (cinq jours pré-opératoires) avant chirurgie digestive lourde chez les malades non dénutris atteints d'un cancer digestif

Comparaison du groupe Impact pré-opératoire au groupe contrôle

Calcul du coût direct des complications, Frais de laboratoires, Traitements, Matériel supplémentaire, Ré-interventions chirurgicales ou autres gestes, Consultations de suivi, Forfait par jour d'hospitalisation

Chirurgie colo-rectale : 288 €/jour

Chirurgie oeso-gastrique : 365 €/jour

Chirurgie pancréatique : 419 €/jour

Forfait de 930 €/jour de soins intensifs

Coût du traitement nutritionnel


0148-6071/05/2901-0S57\$03.00/0

JOURNAL OF PARENTERAL AND ENTERAL NUTRITION

Copyright © 2005 by the American Society for Parenteral and Enteral Nutrition

Preoperative Immunonutrition: Cost-Benefit Analysis

Marco Braga, MD*; and Luca Gianotti, MD, ScD†



Preoperative Immunonutrition: Cost-Benefit Analysis

Marco Braga, MD*; and Luca Gianotti, MD, ScD†

TABLE I *Outcome variables**

	Conventional	Preoperative
Patients with infectious complications	31	14†
Patients with noninfectious complications	36	30
Patients with any complication	49	36
Length of hospital stay (days)	14.0 (7.7)	11.6 (4.7)‡

Preoperative Immunonutrition: Cost-Benefit Analysis

Marco Braga, MD*; and Luca Gianotti, MD, ScD†

TABLE II *Postoperative complications in detail*

	Conventional group	Preoperative group	Total
Anastomotic leak	10	4	14
Abdominal abscess	10	4	14
Pancreatic fistula	5	5	10
Wound infection	11	7	18
Respiratory failure	6	6	12
Delayed gastric emptying	5	6	11
Respiratory tract infection	8	3	11
Sepsis	2	0	2
Wound dehiscence	3	3	6
Circulatory insufficiency	4	6	10
Urinary tract infection	5	4	9
Bleeding	4	2	6
Pleural effusion	4	2	6
Bacteriemia	5	0	5
Intestinal obstruction	1	1	2
Renal dysfunction	2	1	3
Overall	85	54	139

Preoperative Immunonutrition: Cost-Benefit Analysis

Marco Braga, MD*; and Luca Gianotti, MD, ScD†

TABLE VII *Cost-effectiveness analysis*

	Conventional	Preoperative
Total complication costs	315,078	176,282
Complication costs/patient	3089	1728
Nutritional costs/patient	33	144
Total costs/patient	3122	1872*
Effectiveness, %	50.0	62.8
Cost-effectiveness	6244	2985

Preoperative Immunonutrition: Cost-Benefit Analysis

Marco Braga, MD*; and Luca Gianotti, MD, ScD†

TABLE VI *Total costs and DRG reimbursement rates*

	Conventional	Preoperative
Uncomplicated patients costs	184,725	229,208
Complicated patients costs	535,236	334,148
Nutrition costs	3407	14,729
Total costs	723,368	578,085
DRG reimbursement	781,392	740,301

DRG, diagnosis-related group.

Conséquences médico-économiques


0148-6071/05/2901-0857\$03.00/0

JOURNAL OF PARENTERAL AND ENTERAL NUTRITION

Copyright © 2005 by the American Society for Parenteral and Enteral Nutrition

Preoperative Immunonutrition: Cost-Benefit Analysis

Marco Braga, MD*; and Luca Gianotti, MD, ScD†



Encore une étude sur la nutrition...

THE NEW ENGLAND JOURNAL OF MEDICINE

OCCASIONAL NOTES

Chocolate Consumption, Cognition, and Nobel Laureates

Franz H. Messerli

enhanced performance on some cognitive tests, and improved cognitive function in elderly patients with mild impairment have been associated with a regular intake of flavonoids.^{1,2} A subclass of flavonoids called flavanols, which are widely present in cocoa, green tea, red wine, and some fruits, seems to be effective in slowing down or even reversing the reductions in cognitive performance that occur with aging. Dietary flavanols have also been shown to improve endothelial function and to lower blood pressure by causing vasodilation in the peripheral vasculature and in the brain.^{3,4} Improved cognitive performance with the administration of a cocoa polyphenolic extract has even been reported in aged Wistar-Kyoto rats.⁵


Since chocolate consumption could hypothetically improve cognitive function not only in individuals but also in whole populations, I wondered whether there would be a correlation between a country's level of chocolate consumption and its population's cognitive function. To my knowledge, no data on overall national cognitive function are publicly available. Conceivably, however, the total number of Nobel laureates per capita could serve as a surrogate end point reflecting the proportion with superior cognitive function and thereby give us some measure of the overall cognitive function of a given country.

METHODS

A list of countries ranked in terms of Nobel laureates per capita was downloaded from Wikipedia (http://en.wikipedia.org/wiki/List_of_countries_by_Nobel_laureates_per_capita). Be-

Thus, the numbers must of Nobel laureates for every in a given country.

All Nobel Prizes that October 10, 2011, were capita yearly chocolate consumption was obtained from Theobroma-cacao (www.theobroma-cacao.com) and Caobisco (www.caobisco.com). Data were available from (Switzerland), from 2010 to 2004 for 5 countries, and try (China).



There was a close, significant linear correlation ($r=0.791$, $P<0.0001$) between chocolate consumption per capita and the number of Nobel laureates per 10 million persons in a total of 23

Il existe une corrélation linéaire significative ($p<0,0001$) entre la consommation de chocolat par habitant et le nombre de prix Nobel par 10 millions d'habitants sur un total de 23 pays.

Merci pour votre attention

